


8 Week Schedule		 The University of Vermont Health Network - Champlain Valley Physicians Hospital		Mavyret Treatment Worksheet			12 Week Schedule	
Eligibility				Glecaprevir/Pibrentasvir			Eligibility	
Any Genotype				Genotypes 1-6: Treatment Naïve			Any Genotype	
No Cirrhosis				Patient Name:			Cirrhosis (Child Pugh Class A)	
				DOB:			Click here to calculate Child Pugh Score If Child Pugh Class B or C: Do Not Administer - Refer	
8 Week Schedule		Time 0	3 Weeks	9 Weeks	10 Weeks	21 Weeks	22 Weeks	
*Date:								
Recommended Actions:		Order Mavyret 3 pills PO daily with food (84 tablets) x 1 refill Send script to CVPH OutPT Pharmacy	Draw VL & LFTs; Stop treatment if LFTs increase \geq 3x baseline Suggest referral	Draw VL & LFTs; Assess ETR (end of treatment response)	Call patient or schedule f/u to discuss ETR results: VL Undetectable: Continue on to week 21 VL Detectable: Suggest referral	Repeat VL & LFTs; Assess SVR (sustained virological response)	Schedule f/u to discuss SVR results: VL Undetectable: Done - successful SVR VL Detectable: Failed - suggest referral	
Results:	VL	**			XX		XX	
	LFTs	**			XX		XX	
12 Week Schedule		Time 0	3 Weeks	9 Weeks	13 Weeks	14 Weeks	25 Weeks	26 Weeks
*Date:								
Recommended Actions:		Order Mavyret 3 pills PO daily with food; (84 tablets) x 2 refills Send script to CVPH OutPT Pharmacy	Draw VL & LFTs; Stop treatment if LFTs increase \geq 3x baseline Suggest referral	Draw LFTs; Stop treatment if LFTs increase \geq 3x baseline Suggest referral	Draw VL & LFTs; Assess ETR (end of treatment response)	Call patient or schedule f/u to discuss ETR results: VL Undetectable: Continue on to week 25 VL Detectable: Suggest referral	Draw VL & LFTs; Assess SVR (sustained virological response)	Schedule f/u to discuss SVR results: VL Undetectable: Done† - successful SVR VL Detectable: Failed - suggest referral
Results:	VL	**		XX		XX		XX
	LFTs	**				XX		XX
* Dates approximate based on average time for prior approval authorization (average 1 wk)								
** Can be most recent value obtained within the past year								
† Regardless of SVR results, all patients with cirrhosis should be referred to GI for an initial screening EGD (R/O varices) and have life-long serial hepatic ultrasounds and AFP levels performed every 6 months (for hepatoma surveillance). Refer to GI again if either test becomes abnormal.								